

Y1M11 XMQs and MS

(Total: 52 marks)

1. P3_Sample Q6 . 6 marks - Y1M11 Variable acceleration
2. P32(AS)_2018 Q8 . 10 marks - Y1M11 Variable acceleration
3. P32(AS)_2019 Q3 . 8 marks - Y1M11 Variable acceleration
4. P32(AS)_2020 Q3 . 9 marks - Y1M11 Variable acceleration
5. P32(AS)_2021 Q2 . 10 marks - Y1M11 Variable acceleration
6. P32(AS)_2022 Q3 . 9 marks - Y1M11 Variable acceleration

Mathvault.io Solutions

SECTION B: MECHANICS

Answer ALL questions. Write your answers in the spaces provided.

Unless otherwise indicated, whenever a numerical value of g is required, take $g = 9.8 \text{ m s}^{-2}$ and give your answer to either 2 significant figures or 3 significant figures.

6. At time t seconds, where $t \geq 0$, a particle P moves so that its acceleration $\mathbf{a} \text{ m s}^{-2}$ is given by

$$\mathbf{a} = 5t\mathbf{i} - 15t^{\frac{1}{2}}\mathbf{j}$$

When $t = 0$, the velocity of P is $20\mathbf{i} \text{ m s}^{-1}$

Find the speed of P when $t = 4$

(6)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

| Question | Scheme | Marks | AOs |
|--|--|-------|------------------|
| 6 | Integrate \mathbf{a} w.r.t. time | M1 | 1.1a |
| | $\mathbf{v} = \frac{5t^2}{2}\mathbf{i} - 10t^{\frac{3}{2}}\mathbf{j} + \mathbf{C}$ (allow omission of \mathbf{C}) | A1 | 1.1b |
| | $\mathbf{v} = \frac{5t^2}{2}\mathbf{i} - 10t^{\frac{3}{2}}\mathbf{j} + 20\mathbf{i}$ | A1 | 1.1b |
| | When $t = 4$, $\mathbf{v} = 60\mathbf{i} - 80\mathbf{j}$ | M1 | 1.1b |
| | Attempt to find magnitude: $\sqrt{(60^2 + 80^2)}$ | M1 | 3.1a |
| | Speed = 100 m s ⁻¹ | A1ft | 1.1b |
| | | | (6 marks) |
| Notes: | | | |
| <p>1st M1: for integrating \mathbf{a} w.r.t. time (powers of t increasing by 1) 1st A1: for a correct \mathbf{v} expression without \mathbf{C} 2nd A1: for a correct \mathbf{v} expression including \mathbf{C} 2nd M1: for putting $t = 4$ into their \mathbf{v} expression 3rd M1: for finding magnitude of their \mathbf{v} 3rd A1: ft for 100 m s⁻¹, follow through on an incorrect \mathbf{v}</p> | | | |

| Question | Scheme | Marks | AOs |
|--|---|------------|-------------------|
| 8(a) | Multiply out and differentiate wrt to time (or use of product rule i.e. must have two terms with correct structure) | M1 | 1.1a |
| | $v = 2t^3 - 3t^2 + t$ | A1 | 1.1b |
| | $2t^3 - 3t^2 + t = 0$ and solve: $t(2t - 1)(t - 1) = 0$ | DM1 | 1.1b |
| | $t = 0$ or $t = \frac{1}{2}$ or $t = 1$; any two | A1 | 1.1b |
| | All three | A1 | 1.1b |
| | | (5) | |
| (b) | Find x when $t = 0, \frac{1}{2}, 1$ and 2 : $(0, \frac{1}{32}, 0, 2)$ | M1 | 2.1 |
| | Distance = $\frac{1}{32} + \frac{1}{32} + 2$ | M1 | 2.1 |
| | $2\frac{1}{16}$ (m) oe or 2.06 or better | A1 | 1.1b |
| | | (3) | |
| (c) | $x = \frac{1}{2}t^2(t - 1)^2$ | M1 | 3.1a |
| | $\frac{1}{2}$ perfect square so $x \geq 0$ i.e. never negative | A1 cso | 2.4 |
| | | (2) | |
| | | | (10 marks) |
| Notes: | | | |
| (a) | | | |
| M1: Must have 3 terms and at least two powers going down by 1 | | | |
| A1: A correct expression | | | |
| DM1: Dependent on first M, for equating to zero and attempting to solve a <u>cubic</u> | | | |
| A1: Any two of the three values (Two correct answers can imply a correct method) | | | |
| A1: The third value | | | |
| (b) | | | |
| M1: For attempting to find the values of x (at least two) at their t values found in (a) or at $t = 2$ or equivalent e.g. they may integrate their v and sub in at least two of their t values | | | |
| M1: Using a correct strategy to combine their distances (must have at least 3 distances) | | | |

A1: $2\frac{1}{16}$ (m) oe or 2.06 or better

(c)

M1: Identify strategy to solve the problem such as:

- (i) writing x as $\frac{1}{2} \times$ perfect square
- (ii) or using x values identified in (b).
- (iii) or using calculus i.e. identifying min points on $x-t$ graph.
- (iv) or using $x-t$ graph.

A1 cso : Fully correct explanation to show that $x \geq 0$ i.e. never negative

Mathvault.io Solutions

3. A particle, P , moves along a straight line such that at time t seconds, $t \geq 0$, the velocity of P , $v \text{ ms}^{-1}$, is modelled as

$$v = 12 + 4t - t^2$$

Find

- (a) the magnitude of the acceleration of P when P is at instantaneous rest, (5)
- (b) the distance travelled by P in the interval $0 \leq t \leq 3$ (3)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



| Question | Scheme | Marks | AOs | Notes |
|-------------|--|------------------|------|---|
| 3(a) | $v = 12 + 4t - t^2 = 0$ and solving | M1 | 3.1a | Equating v to 0 and solving the quadratic If no evidence of solving, and at least one answer wrong, M0 |
| | $t = 6$ (or -2) | A1 | 1.1b | 6 but allow -2 as well at this stage |
| | Differentiate v wrt t | M1 | 1.1a | For differentiation (both powers decreasing by 1) |
| | $(a = \frac{dv}{dt} \Rightarrow) 4 - 2t$ | A1 | 1.1b | Cao; only need RHS |
| | When $t = 6$, $a = -8$; Magnitude is $8 \text{ (m s}^{-2}\text{)}$ | A1 | 1.1b | Substitute in $t = 6$ and get $8 \text{ (m s}^{-2}\text{)}$ as the answer . Must be positive . (A0 if two answers given) |
| | | (5) | | |
| (b) | Integrate v wrt t | M1 | 3.1a | For integration (at least two powers increasing by 1) |
| | $(s \Rightarrow) 12t + 2t^2 - \frac{1}{3}t^3 (+C)$ | A1 | 1.1b | Correct expression (ignore C) only need RHS Must be used in part (b) |
| | $t = 3 \Rightarrow \text{distance} = 45 \text{ (m)}$ | A1 | 1.1b | Correct distance. Ignore units |
| | | (3) | | |
| | | (8 marks) | | |

| Question | Scheme | | Marks | AOs |
|------------------|--|--|-------|------|
| 3(a) | $v = 3t - 2t^2 + 14$ and differentiate | | M1 | 3.1a |
| | $a = \frac{dv}{dt} = 3 - 4t$ or $(7 - 2t) - 2(t + 2)$ using product rule | | A1 | 1.1b |
| | $3 - 4t = 0$ and solve for t | | M1 | 1.1b |
| | $t = \frac{3}{4}$ oe | | A1 | 1.1b |
| | | | (4) | |
| 3(b) | Solve problem using $v = 0$ to find a value of t $\left(t = \frac{7}{2}\right)$ | | M1 | 3.1a |
| | $v = 3t - 2t^2 + 14$ and integrate | | M1 | 1.1b |
| | $s = \frac{3t^2}{2} - \frac{2t^3}{3} + 14t$ | | A1 | 1.1b |
| | Substitute $t = \frac{7}{2}$ into their s expression (M0 if using <i>suvat</i>) | | M1 | 1.1b |
| | $s = \frac{931}{24} = 38\frac{19}{24} = 38.79166..(m)$ Accept 39 or better | | A1 | 1.1b |
| | | | (5) | |
| (9 marks) | | | | |
| Notes: | | | | |
| (a) | M1 | Multiply out and attempt to differentiate, with at least one power decreasing | | |
| | A1 | Correct expression | | |
| | M1 | Equate their a to 0 and solve for t | | |
| | A1 | cao | | |
| (b) | M1 | Uses $v = 0$ to obtain a value of t | | |
| | M1 | Attempt to integrate, with at least one power increasing | | |
| | A1 | Correct expression | | |
| | M1 | Substitute in their value of t , which must have come from using $v = 0$, into their s (must have integrated) | | |
| | A1 | 39 or better | | |

| Question | Scheme | | Marks | AOs |
|-------------------|---|--|-------|------|
| 2(a) | Differentiate v w.r.t. t | | M1 | 3.1a |
| | $a = \frac{dv}{dt} = 10 - 2t$ isw | | A1 | 1.1b |
| | | | (2) | |
| 2(b) | Solve problem using $v = 0$ when $t = 6$ | | M1 | 3.1a |
| | $0 = 10t - t^2 - 24$ | | A1 | 1.1b |
| | Solve quadratic oe to find other value of t | | M1 | 1.1b |
| | $t = 4$ | | A1 | 1.1b |
| | | | (4) | |
| 2(c) | Integrate v or $-v$ w.r.t. t | | M1 | 3.1a |
| | $5t^2 - \frac{1}{3}t^3 - 24t$ | | A1 | 1.1b |
| | Total distance = $-\left[5t^2 - \frac{1}{3}t^3 - 24t\right]_0^4 + \left[5t^2 - \frac{1}{3}t^3 - 24t\right]_4^6$ | | M1 | 2.1 |
| | $\frac{116}{3}$ (m) | | A1 | 1.1b |
| | | | (4) | |
| (10 marks) | | | | |
| Notes: | | | | |
| 2a | M1 | Differentiate, with both powers decreasing by 1 | | |
| | A1 | Correct expression | | |
| 2b | M1 | Put $t = 6$ OR use $(t-6)(t-x) = t^2 - 10t + k$ oe | | |
| | A1 | Correct expression (unsimplified) for v OR $v = (t-6)(t-4)$ | | |
| | M1 | Put $v = 0$ to give quadratic in t and solve for other value of t | | |
| | A1 | $t = 4$ | | |
| 2c | M1 | Integrate, with at least two powers increasing by 1 (allow if only two terms integrated) | | |
| | A1 | Correct expression | | |
| | M1 | Complete method to find the total distance | | |
| | A1 | Accept 39(m) or better | | |

3. A fixed point O lies on a straight line.

A particle P moves along the straight line.

At time t seconds, $t \geq 0$, the distance, s metres, of P from O is given by

$$s = \frac{1}{3}t^3 - \frac{5}{2}t^2 + 6t$$

- (a) Find the acceleration of P at each of the times when P is at instantaneous rest. (6)
- (b) Find the total distance travelled by P in the interval $0 \leq t \leq 4$ (3)



| Question | Scheme | Marks | AOs |
|------------------|---|--|------------|
| 3(a) | Differentiate s wrt t | M1 | 3.1a |
| | $(v \Rightarrow) t^2 - 5t + 6$ | A1 | 1.1b |
| | Equate their v to 0 and solve | M1 | 1.1b |
| | $t = 2$ or 3 | A1 | 1.1b |
| | $(a \Rightarrow) 2t - 5$ | B1ft | 2.1 |
| | $a = 1$ and -1 (m s^{-2}) isw (A0 if extras) | A1 | 1.1b |
| | | (6) | |
| (b) | Attempt to find values of s for $t = 2, 3$ and 4 oe Correct values are $\left(s_2 = \frac{14}{3}, s_3 = \frac{9}{2} \text{ and } s_4 = \frac{16}{3} \right)$ Could be implied by correct values for: $s_2, (s_3 - s_2)$ and $(s_4 - s_3)$ which are $\frac{14}{3}, \left(-\frac{1}{6}\right)$ and $\frac{5}{6}$ | DM1 | 1.1b |
| | Total distance travelled $= s_2 + (s_2 - s_3) + s_4 - s_3$ OR $s_2 - (s_3 - s_2) + s_4 - s_3$ OR $\left[\frac{1}{3}t^3 - \frac{5}{2}t^2 + 6t \right]_0^2 - \left[\frac{1}{3}t^3 - \frac{5}{2}t^2 + 6t \right]_2^3 + \left[\frac{1}{3}t^3 - \frac{5}{2}t^2 + 6t \right]_3^4$ OR $\frac{14}{3} - \left(-\frac{1}{6}\right) + \frac{5}{6}$ OR $s_2 + 2(s_2 - s_3) + s_4 - s_2$ $(= 2s_2 - 2s_3 + s_4)$ oe | M1 | 2.1 |
| | $5\frac{2}{3}$ oe (m) Accept 5.7 or better | A1 | 1.1b |
| | | | (3) |
| (9 marks) | | | |
| Notes: | | | |
| 3a | M1 | Differentiate, with at least 2 powers decreasing by 1 | |
| | A1 | Correct expression | |
| | M1 | Must have attempted to differentiate s to find v and be solving a 3 term quadratic | |
| | A1 | Both values needed | |
| | B1ft | Follow their v (must be differentiating) | |

| | | |
|-----------|----------------|--|
| | A1 | cao |
| 3b | DM 1 | This mark is dependent on the 2 nd M1 in part (a) and their t values are between 0 and 4. Clear attempt to find all three s values (may integrate their v incorrectly) N.B. No penalty for extra values. |
| | M1 | Complete method using their s values Do NOT condone sign errors. |
| | A1 | Any equivalent fraction, 5.7 or better. |
| | | S.C. Correct answer, with no working, scores all 3 marks, since $\int_0^4 t^2 - 5t + 6 dt$ entered on a calculator will give $\frac{17}{3}$ |

Mathvault.io Solutions